

Table supplement3. Patients' blood analysis results obtained immediately after surgical operation (a/o). Median values and minimum-maximum values are presented.

	IL-8, pg/ml	IL-10, pg/ml	IL-6, pg/ml	MPO, ng/ml	Thiols, 10 ⁻⁶ M	CL, V
Group 1	3,4 (0,1÷6,4)	6,0* (1,3÷10,7)	13,6 (0,7÷110)	39,8* (23÷312)	185* (123÷280)	13* (7÷25)
Group 2	1,1 (0,9÷10,4)	6,1* (3,5÷56,4)	40,7*, ^Y (0,6÷288)	35,1 * (11,7÷54,0)	232 (189÷270)	14*, ¹ (12÷26)
Group 3	23,9 *, ^{1,2} (0,2÷430)	36,5 *, ¹ (1,7÷132)	375*, ^{1,2} (50÷1754)	87,0*, ^{1,2} (34,5÷342)	128* (13÷232)	14* (0÷53)
Normal values	1,4 (0,7÷1,4)	2,4 (0,2÷11,3)	8,6 (0,1÷21,4)	19,6 (13,5÷40,5)	251 (175÷332)	2,5 (1,6÷5,7)

Statistical differences between groups (Mann-Whitney U-test)

*p<0,05 vs normal values

¹p<0,05 vs group 1

²p<0,05 vs group 2

Statistical differences between "b/o" and "a/o" values (Wilcoxon paired test)

^Y p<0,05 vs «b/o» value